

# Inquiring Minds Topic – 8 January 2021

Roger Palms, Moderator

## The Politics of Gender

**Local lawmakers seek to ban medical interventions for transgender children**

### **Thought starters:**

1. Is there a danger that children may be making decisions about themselves before they are mature enough to know what they are doing?
2. How is it happening that “blockers” rather than counseling seem to be the first choice of parents and doctors?
3. Children are not always kind. How will a child’s mental health be affected if he/she shows up at school as the other gender?
4. What help/guidance needs to be offered to parents who feel torn about their child’s gender dysphoria?
5. At what point should lawmakers be getting involved?

James Younger’s mother believed he, James, should wear dresses and grow his hair long because he considered himself a girl. His father claimed the seven-year-old was perfectly content with trousers, short hair and being a boy. Almost every detail of the couple’s vicious custody battle was fought over their child’s gender identity. After his mother won, culture warriors piled in. Ted Cruz, a senator from Texas, said the child was “a pawn in a left-wing political agenda.”

Lawmakers in a number of states say the Youngers’ row has prompted them to try to pass laws banning medical interventions that bring transgender children closer to the sex with which they identify. At least five states have drawn up bills which would make it illegal to perform gender-re-assignment surgery on children and to prescribe puberty blockers and cross-sex hormones for them.

The danger of making trans rights an issue in the culture wars is that it prevents a discussion of the dangers of prescribing blockers and sex hormones for children who suffer from

gender dysphoria, the distress caused by feeling that one's sex at birth and gender identity do not match.

Data on all aspects of transgender medical interventions are poor. No one knows how many children have been prescribed these drugs. Little is known about how they have fared since. But in the past decade there has been a surge in the number of children treated as trans. Clinics serving them have mushroomed. In 2007 there was one. Today there are perhaps 50. Waiting lists at many are long and lengthening.

Anecdotal evidence suggests that standards of care have failed to keep pace. The biggest concern is that children put on blockers—first prescribed between the ages of 9 and 14 to suppress the action of sex hormones—and later, testosterone or estrogen, do not first undergo sufficiently comprehensive evaluations. Guidelines from the World Professional Association for Transgender Health say such interventions should follow “extensive exploration of psychological, family and social issues.” That seems elementary. There is no medical test for gender dysphoria. Research suggests that most children who identify as the other sex eventually grow out of it. They are also more likely to suffer from anxiety and depression. Untangling all of this and establishing whether a child is likely to go on feeling that they are in the wrong body—a guess, at best—poses significant challenges for children, parents and their doctors.

Laura Edwards-Leeper, a professor of psychology at Pacific University in Oregon, who helped found America's first transgender clinic for children in Boston, reckons the “vast majority” of children on blockers or sex hormones have not undergone proper assessments. This, she says, is because of a shortage of mental-health professionals with the necessary training and the desire of doctors to provide care for a group that has long been denied it.

This carries the obvious risk that patients will regret transitioning. No one knows how many people fall into this category. A small number of those who were put on blockers and sex hormones have since “detransitioned.” The most outspoken among them are lesbians who

say that had they been encouraged to explore gender non-conformity—the idea, for instance, that women can be butch—rather than transgenderism, they would not have taken testosterone. Others say mental-health problems caused their gender dysphoria and cross-sex hormones were prescribed as the solution.

A second, related problem concerns the way blockers are sold to patients and their families. Developed in the 1980s to treat premature puberty, they have transformed transgender health care since they were first used for this purpose in the late 1990s. Doctors attest that they save adolescents who feel desperate about developing the “wrong” sex characteristics from enormous distress. Blockers can forestall more traumatic interventions later: the removal of breasts, or the shaving of an Adam’s apple. Their effects are largely reversible. Doctors who prescribe them routinely refer to blockers as a “conservative” measure.

Yet few children seem to step off the treatment path that blockers set them on. The great majority go on to sex hormones. Given the inadequacy of many pre-treatment evaluations, this seems unlikely to be wholly the result of sound diagnoses.

Puberty blockers also have other side effects. Over time, they can affect bone density. This means that doctors are keen to move patients who want to continue treatment onto sex hormones within a few years. But many of the effects of these are irreversible, including infertility.

Paul Hruz, an endocrinologist at Washington University School of Medicine in St Louis, says interrupting puberty may have other harmful effects. A surge of hormones during puberty may help put adolescents at ease with their birth gender. Puberty blockers would prevent that process.

Few doctors worried by these problems are prepared to speak about them openly. That is unsurprising given how inflammatory the issue has become. When Lisa Littman, a professor of behavioral and social sciences at Brown University, published a paper in 2018 in which she noted that most transgender children were teenage girls with no history of gender

dysphoria—a phenomenon she called “rapid-onset gender dysphoria” —she was denounced as transphobic.

In such a polarized environment, bills proposing blanket bans of puberty blockers are likely to be counterproductive. They may push advocates for early intervention to further extremes. A better approach would be twofold. A neutral assessment of the existing data on the use of blockers, hormones and their effects would help patients and their families make decisions. Most existing research has been undertaken by those working in the field. At the same time, clinics should ensure that children in transgender clinics undergo comprehensive mental-health evaluations. For all this to happen there needs to be an acknowledgment of the dangers of starting children on often-irreversible treatments.

Taken from *The Economist* February, 2020

### **From the Mayo Clinic**

## **How can you support a gender-nonconforming or transgender child?**

If your child is persistent about gender identity feelings, listen. Talk to your child and ask questions without judgment. To support your child:

- Don't assume your child's gender expression is a form of rebellion or defiance.
- Don't prevent your child from expressing gender in public or at family activities to avoid it making you or someone else uncomfortable.
- Don't try to shame or punish the gender expression out of your child.
- Don't block your child's access to gender-diverse friends, activities or resources.
- Don't blame your child for experiencing discrimination.
- Don't belittle or ridicule your child's gender expression or allow others in your family to do so.

Speak positively about your child to your child and to others. Show your admiration for your child's identity and expression of it. By allowing your child to demonstrate preferences and share them, you'll encourage a positive sense of self and keep the lines of communication open.

Also, try to let go of specific fantasies you might have had about your child's future and, instead, focus on what brings your child joy and security. A child living with supportive parents and caregivers is likely to be a happier child.